© 1999 Nature America Inc. • http://medicine.nature.com

CORRECTION

Modulation of oncogenic potential by alternative gene use in human prostate cancer

S.S. KADKOL, J.R. BRODY, J. PEVSNER, J.BAI & G.R. PASTERNACK

*Nature Med.***5**, 275–279 (1999).

On page 278, the original Fig. 3, ommitted four gap symbols and two amino-acid changes. The correct figure is shown.

```
          1                                                  50
pp32      memgrrihle lrnrtpsdvk elvldnsrsn egklegltde feeleflsti
pp32r1          s      a                    a             k
pp32r2        kw                  f    q                  l n

          51                                                 100
pp32      nvgltsianl pklnklkkle lsdnrvsggl evlaekcpnl thlnlsgnki
pp32r1     g     sd     ~  r     ~~~k                   y
pp32r2     i                     s  a v              i

          101                                                150
pp32      kdlstieplk klenlksldl fncevtnlnd yrenvfkllp qltyldgydr
pp32r1              q                     g       l       scyw
pp32r2                   e      t       n  ~~~~~~~~~~ ~~~~~~~~~~

          151                                                200
hpp32p    ddkeapdsda egyvegldde eededeeeyd edaqvvedee dedeeeegee
pp32r1     h    y  i  dh           g h                 g e
pp32r2    ~~~~~~~~~~ ~~~~~~~~~~ ~~~~~~~~~~ ~~~~~~~~~~ ~~~~~~~~~~

          201                                                249
hpp32p    edvsgeeeed eegyndgevd deedeeelge eerggkrkre pedegeddd
pp32r1         gd              g                  ~~ ~~~~~~~~~
pp32r2    ~~~~~~~~~~ ~~~~~~~~~~ ~~~~~~~~~~ ~~~~~~~~~~ ~~~~~~~~~
```

ERRATUM

Mice deficient in tumor necrosis factor-α are resistant to skin carcinogenesis

R.J. MOORE, D.M. OWENS, G. STAMP, C. ARNOTT, F. BURKE, N. EAST, H. HOLDSWORTH, L.TURNER, B. ROLLINS, M. PASPARAKIS, G. KOLLIAS & F. BALKWILL

Nature Med. **5**, 828–831 (1999).

On page 829, the legend for Fig 1 was incorrect. The corrected figure and legend are shown here. We regret this error.



Fig. 1 Skin carcinogenesis in wild-type and TNF-$\alpha^{-/-}$ mice. ***a***, Initiation promotion. Mice were treated with one dose of 25 μg DMBA and then 4 μg TPA three times weekly for 15 weeks. C57Bl6/129 (■), $n = 14$; TNF-$\alpha^{-/-}$ (●), $n = 15$. ***b***, Initiation promotion. Mice were treated with one dose of 25 μg DMBA and then 4 μg TPA twice weekly for 15 weeks. Balb/c (■), $n = 14$; TNF-$\alpha^{-/-}$ (●), $n = 13$. ***c***, Complete carcinogenesis. Mice were treated with 5.1 μg DMBA twice weekly for 20 weeks. C57Bl6/129 (■), $n = 15$; TNF-$\alpha^{-/-}$ (●), $n = 20$. ***d***, Initiation promotion. Mice were treated with one dose of 25 μg DMBA and then 4 μg TPA twice weekly. C57Bl6/129 (■), $n = 15$; MCP-$1^{-/-}$ (●), $n = 15$.

© 1999 Nature America Inc. • http://medicine.nature.com